

Epstein-Barr Virus-Associated Hemophagocytic Syndrome and Fatal Infectious Mononucleosis

Motohiko Okano and Thomas G. Gross

Department of Pediatrics, Hokkaido University School of Medicine, Sapporo, Japan (M.O.); Departments of Pathology and Microbiology, and Pediatrics, University of Nebraska Medical Center, Omaha, Nebraska (T.G.G.)

Virus-associated hemophagocytic syndrome (VAHS) has been thought to be a distinct clinical entity, characterized by intermittent fever, enlarged liver and spleen, and the appearance of hemophagocytosis. Hemopoietic cells are actively ingested by monocytes/macrophages in various organs, including lymph nodes, bone marrow, liver, and spleen. Epstein-Barr virus (EBV) is now thought to be one of the major causes for the development of this unique syndrome. Additionally, VAHS is often associated with fatal infectious mononucleosis (IM). The relationship between EBV-associated VAHS and fatal IM is discussed in this concise review. © 1996 Wiley-Liss, Inc.

Key words: Epstein-Barr virus, hemophagocytic syndrome, fatal infectious mononucleosis

INTRODUCTION

Hemophagocytic syndromes (HPS) may be divided into different clinical categories, consisting of 1) a sporadic disorder, 2) a form associated with acute infection, 3) a familial form seen in children, and 4) a form associated with malignant disorders, immunodeficiencies, or defective leukocyte function [1]. Histopathologically, lesions are characterized by a mononuclear cell infiltration with a prominent histiocytic proliferation and phagocytosis of blood cells.

Most patients with infection-related HPS are afflicted with viral infection, defining the so-called virus-associated hemophagocytic syndrome (VAHS) [2]. Some investigators have proposed that the clinical syndrome represents a direct response to a fulminant viral infection, whereas others propose that virus(es) triggers an uncontrolled histiocytic proliferation in individuals with an underlying cellular immunoregulatory imbalance. However, precise pathogenetic mechanisms are still undetermined. Recently patients with VAHS have been reported more frequently, and Epstein-Barr virus (EBV) infection is considered to be the cause of most cases. Generally, infectious mononucleosis (IM) is a benign, self-limiting disease associated with a primary EBV infection. However, rare cases develop severe or fatal IM; additionally, severe or fatal IM is commonly accompanied by VAHS [3]. These disorders are the subjects of this report.

HISTORY OF VIRUS-ASSOCIATED HEMOPHAGOCYTIC SYNDROME AND FATAL INFECTIOUS MONONUCLEOSIS

In the late 1970's, Risdall et al. [2] first reported 19 cases characterized by similar clinical symptoms and laboratory findings, consisting of intermittent fever, lymph-node enlargement, hepatosplenomegaly, a tendency to pancytopenia and a hemophagocytosis in various tissues, where hemopoietic cells were actively ingested by monocytes/macrophages. They named this disorder virus-associated hemophagocytic syndrome (VAHS), because most patients had a preceding virus-associated illness caused by herpes or adenoviruses. Additionally, spontaneous recovery was common in many cases. In the early 1980's, cases of VAHS associated with EBV infection were reported, some being fatal [4–6]. In these cases, the presence of IgM antibodies to viral capsid antigen (VCA) and the absence of IgG antibodies to EBV-determined nuclear antigen (EBNA) indicated primary EBV infection. Following these reports, there were additional reports of EBV-associated VAHS [7–11]. Initially, the outcome of

Received for publication March 27, 1996; accepted April 24, 1996.

Address reprint requests to Motohiko Okano, M.D., Ph.D., Department of Pediatrics, Hokkaido University School of Medicine, North 15 West 7, Kita-ku, Sapporo 060, Japan.

EBV-associated VAHS was considered to be favorable. However, with more cases, the prognosis of EBV-associated VAHS was demonstrated to be poor. Some patients exhibited a rapidly deteriorating and fatal course, similar to the clinical course of patients with fatal IM [12,13].

Males with X-linked lymphoproliferative syndrome (XLP) often develop severe or fatal IM accompanied by HPS observed in affected tissues, such as lymph nodes, bone marrow, liver, and spleen [14,15]. Approximately 60% of patients with XLP manifest severe or fatal IM, with VAHS generally associated with primary EBV infection [16]. VAHS associated with XLP and VAHS without XLP are believed to be different clinical entities. However, some of the pathogenetic mechanism(s) may be similar [17]. The study of underlying pathogenetic mechanisms of XLP may clarify the relationship between VAHS and fatal IM.

IMMUNE EVENTS IN PATIENTS WITH EBV-ASSOCIATED VAHS AND FATAL IM

During the acute phase of IM, atypical lymphocytes in blood increase remarkably [18]. Enlarged lymph nodes contain infiltrating CD3-, CD4-, CD8-, CD20-, and NKH-1-positive cells [19]. In the course of the disease, EBV-specific cytotoxic T lymphocytes (CTL), mainly derived from CD8-positive cells, emerge and kill EBV-infected cells [18,20]. In contrast to the increased number of circulating CD8-positive cells, subsets of CD4-positive cells increase slightly, but are important for the immune events responsible for cell-mediated immunity or humoral immunity, since they produce interferon (IFN)-gamma or interleukin (IL)-4 [21]. Severely defective or imbalanced immune defenses against primary EBV infection often lead to fatal IM, primarily due to fulminant hepatitis and VAHS [22].

Recently, studies to ascertain the pathogenetic mechanism(s) of VAHS and fatal IM have focused on a possible imbalance of various cytokine networks [23]. We studied 5 patients with IM during the acute phase, and none showed increased serum levels of IFN-gamma or IFN-alpha [24]. A part of sera from 5 patients with fatal IM demonstrated increased levels of these IFNs. All of these patients succumbed to VAHS. This observation suggested that an imbalance of IFN production was associated with this complication. Markedly elevated levels of soluble IL-2 receptor were also noted in patients with VAHS [25]. Additionally, increased serum levels of IFN-gamma, macrophage-colony stimulating factor (MCSF), tumor necrosis factor (TNF)-alpha, and IL-6 were recently reported in patients with hemophagocytic lymphohistiocytosis [26]. MCSF is produced by T cells, and stimulates monocytes/macrophages. Furthermore, the major source of IL-6 and TNF-alpha is activated monocytes/macrophages. Therefore, an overproliferation of activated T




Fig. 1. Schematic hypotheses of immune events for Epstein-Barr virus (EBV)-associated hemophagocytic syndrome and fatal infectious mononucleosis (IM). EBV infects B cells, and/or T cells, and/or monocytes/macrophages, initially. Following these events, monocytes/macrophages are activated, and ingest various types of hemopoietic cells. Whether EBV directly infects T cells and/or monocytes/macrophages remains to be fully determined. In the case of virus-associated hemophagocytic syndrome or fatal IM, an overproliferation of activated T cells appears to play a significant role in the pathogenesis of the disease, because cytokines which were mainly produced by T cells were elevated in the circulation. sIL-2R, soluble interleukin-2 receptor; MCSF, macrophage-colony stimulating factor; IFN-gamma, interferon-gamma; IL-6, interleukin-6; TNF-alpha, tumor necrosis factor-alpha.

cells appears to be part of the syndrome of HPS, and the activation and/or overproliferation of T cells directed against EBV-infected cells may be a factor in HPS pathogenesis. Using molecular hybridization techniques, some have demonstrated T cells or monocytes/macrophages to be infected with EBV [27–29]. Therefore, EBV may infect T cells or monocytes/macrophages directly, and result in T cell or monocyte/macrophage proliferation. Additionally, NK cells contained EBV genome in one patient with VAHS [30], but these findings require confirmation. Though only B cells have been infected with EBV in healthy seropositive individuals [31], it is possible that other cells may be diseased in actively infected or immunocompromised individuals. Whether the immunophenotype of EBV-infected cells plays a major role in the pathogenesis of EBV-associated VAHS and fatal IM needs further study. Schematic hypotheses for the development of VAHS and fatal IM are shown in Figure 1.

In patients with primary or secondary immunodeficiencies, EBV-induced lymphoproliferative disorders (LPD) occur at an increased frequency [3]. Other than XLP, EBV-associated VAHS has rarely been documented in these patients, although non-EBV-associated VAHS

TABLE I. Therapeutic Approaches for Epstein-Barr Virus (EBV)-Associated Hemophagocytic Syndrome and Fatal Infectious Mononucleosis

Approach	Major target of action
1. Acyclovir	EBV
2. Ganciclovir	EBV
3. Adenine arabinoside	EBV
4. Immunoglobulins	EBV, B cell, T cell
5. Interferon-alpha	EBV, T cell, NK cell
6. Interferon-gamma	EBV, T cell, NK cell
7. Corticosteroids	Monocyte/macrophage, T cell, NK cell
8. Etoposide (VP16)	Monocyte/macrophage
9. Cyclosporine A	T cell
10. Anti-B-cell monoclonal antibodies	B cell
11. Bone marrow transplantation	Stem cell

occurs in patients with either primary or acquired immunodeficiencies. These LPD result from EBV-infected B-cell proliferations. Therefore, EBV-associated VAHS may occur in individuals who were previously immunocompetent. Though approximately 10% of patients with XLP have manifestations of XLP prior to EBV infection (e.g., lymphoma or dysgammaglobulinemia), the majority of patients have no evidence of immunodeficiency prior to EBV infection [16].

We recently described an unusual EBV infection in an XLP patient with fatal IM and VAHS [32]. A spontaneously-established cell line from an enlarged cervical lymph node demonstrated latent EBV infection in which no EBV replication occurred. In contrast, a combination of latent and lytic infection, in which EBV replication could occur, was noted in cell lines from blood lymphocytes. Latently infected cells may be less susceptible to lysis by natural killer (NK) cells and EBV-specific CTL. Restriction endonuclease enzyme-digested patterns of EBV were similar at each site, confirming one strain of virus, despite differing states of infection. These observations suggest that unknown defective controlling mechanisms for EBV infection exist at cellular levels in patients with severe EBV infection. Additionally, a variant EBV strain was isolated in 2 siblings with fatal lymphoproliferative disease [33]. This evidence indicated that a variant EBV strain might be responsible at least in part for the fatal course of these siblings, although further investigations are needed.

THERAPEUTIC APPROACHES FOR EBV-ASSOCIATED VAHS AND FATAL IM

Many therapies have been attempted against severe EBV infection [3]. These treatments include antiviral agents (such as acyclovir, ganciclovir, and adenine arabinoside), immunoglobulins, and immunomodulating agents such as IL-2, IFN-alpha, IFN-gamma, and corticosteroids. However, no beneficial effects have been clearly demonstrated from any of these treatments. Recently, eto-

poside was given to a patient with severe IM accompanied by VAHS who had XLP resulting in a clinical remission [34]. Etoposide mainly reduces the activity of monocytes/macrophages in vitro [35]. Although further clinical trials are necessary, Chinese investigators recently described an immunomodulatory regimen of intravenous immunoglobulins and/or etoposide that improved the prognosis of patients with HPS [36]. Additionally, cyclosporine A (CSA) or tacrolimus (formerly named FK506) may be useful in controlling the proliferation of T cells in spite of decreasing activity of EBV-specific CTL [3,37]. Of the over 150 affected males in the XLP registry (University of Nebraska Medical Center) who have had severe or fatal IM, there are 5 long-term survivors; 4 boys received etoposide and immunosuppressive therapies such as CSA (unpublished observations). Taken together, the current recommended approach for the treatment of VAHS or fatal IM is the use of etoposide with or without an immunomodulatory regimen. However, anti-B-cell monoclonal antibodies may be considered when infected B cells are thought to be a major factor in pathogenesis [38]. When underlying genetic factors for controlling EBV infection exist, such as XLP or familial HPS, intensive therapy supported by bone-marrow transplantation may be the treatment of choice [16,34]. Therapeutic approaches against EBV-associated VAHS and fatal IM, and a major target of action, are summarized in Table I.

CONCLUSIONS

In the 30 years since the discovery of EBV, this virus has been etiologically linked to a spectrum of human diseases. Previously unknown relationships may yet be detected by new diagnostic procedures. Furthermore, little is known about the pathogenetic mechanisms of EBV-associated diseases. Further advances in the recognition and differentiation of various EBV-associated entities should provide a more rational basis for the assessment of potentially effective therapies.

ACKNOWLEDGMENTS

The authors are indebted to their colleagues at the Department of Pathology and Microbiology, and the Department of Pediatrics, University of Nebraska Medical Center, Omaha, Nebraska, and at the Department of Virology, Cancer Institute, Hokkaido University School of Medicine and the Department of Pediatrics, Hokkaido University School of Medicine, Sapporo, Japan, for their helpful discussions and comments. This work was partly supported by a Grant in Aid for Cancer Research from the Foundation for Promotion of Cancer Research, Japan, and by the William C. Havens Foundation, Omaha, Nebraska. Finally, this work is dedicated to the memory of David T. Purtilo, M.D., who was a pioneer in the study of Epstein-Barr virus infection and immunodeficiencies.

REFERENCES

- Favara BE: Hemophagocytic lymphohistiocytosis: A hemophagocytic syndrome. *Semin Diagn Pathol* 9:63, 1992.
- Risdall RJ, McKenna RW, Nesbit ME, Krivit W, Balfour HH Jr, Simmons RL, Brunning RD: Virus-associated hemophagocytic syndrome. A benign histiocytic proliferation distinct from malignant histiocytosis. *Cancer* 44:993, 1979.
- Okano M, Thiele GM, Davis JR, Grierson HL, Purtilo DT: Epstein-Barr virus and human diseases: Recent advances in diagnosis. *Clin Microbiol Rev* 1:300, 1988.
- Wilson ER, Malluk A, Stagno S, Crist WM: Fatal Epstein-Barr virus-associated hemophagocytic syndrome. *J Pediatr* 98:260, 1981.
- Reisman RP, Greco MA: Virus-associated hemophagocytic syndrome due to Epstein-Barr virus. *Hum Pathol* 15:290, 1984.
- Sullivan JL, Woda BA, Herrod HG, Koh G, Rivara FP, Mulder C: Epstein-Barr virus-associated hemophagocytic syndrome: Virological and immunopathological studies. *Blood* 65:1097, 1985.
- McClain K, Gehrz R, Grierson H, Purtilo D, Filipovich A: Virus-associated histiocytic proliferations in children. Frequent association with Epstein-Barr virus and congenital or acquired immunodeficiencies. *Am J Pediatr Hematol Oncol* 10:196, 1988.
- Oshima K, Kikuchi M, Eguchi F, Kobari S, Tasaka H: Virus-associated haemophagocytic syndrome with Epstein-Barr virus infection. *Virchows Arch [A]* 419:519, 1991.
- Dreyer ZE, Dowell BL, Chen H, Hawkins E, McClain KL: Infection-associated hemophagocytic syndrome. Evidence for Epstein-Barr virus gene expression. *Am J Pediatr Hematol Oncol* 13:476, 1991.
- Kawaguchi H, Miyashita T, Herbst H, Niedobitek G, Asada M, Tsuchida M, Hanada R, Kinoshita A, Sakurai M, Kobayashi N, Mizutani S: Epstein-Barr virus-infected T lymphocytes in Epstein-Barr virus-associated hemophagocytic syndrome. *J Clin Invest* 92:1444, 1993.
- Kikuta H, Sakiyama Y, Matsumoto S, Oh-Ishi T, Nakano T, Nagashima T, Oka T, Hironaka T, Hirai K: Fatal Epstein-Barr virus-associated hemophagocytic syndrome. *Blood* 82:3259, 1993.
- Penman HG: Fatal infectious mononucleosis. A critical review. *J Clin Pathol* 23:756, 1970.
- Harries JT, Ferguson AW: Fatal infectious mononucleosis with liver failure in two sisters. *Arch Dis Child* 43:480, 1968.
- Purtilo DT, Cassel CK, Yang JPS, Harper K, Stephenson SR, Landing BH, Vawter GF: X-linked recessive progressive combined variable immunodeficiency (Duncan's disease). *Lancet* i:935, 1975.
- Purtilo DT, Grierson HL, Davis JR, Okano M: The X-linked lymphoproliferative disease: From autopsy toward cloning the gene 1975-1990. *Pediatr Pathol* 11:685, 1991.
- Seemayer TA, Gross TG, Egeler RM, Pirruccello SJ, Davis JR, Kelley CM, Okano M, Lanyi A, Sumegi Y: X-linked lymphoproliferative disease: Twenty-five years after discovery. *Pediatr Res* 38:471, 1995.
- Mroczek EC, Weisenburger DD, Grierson HL, Markin R, Purtilo DT: Fatal infectious mononucleosis and virus-associated hemophagocytic syndrome. *Arch Pathol Lab Med* 111:530, 1987.
- De Waele M, Thielemans C, Van Camp BKG: Characterization of immunoregulatory T cells in EBV-induced infectious mononucleosis by monoclonal antibodies. *N Engl J Med* 304:460, 1981.
- Prange E, Trautmann JC, Kreipe H, Radzun HZ, Parwareschi M: Detection of Epstein-Barr virus in lymphoid tissues of patients with infectious mononucleosis by in situ hybridization. *J Pathol* 166:133, 1992.
- Royston I, Sullivan JL, Periman PO, Perlin E: Cell-mediated immunity to Epstein-Barr virus transformed lymphoblastoid cells in acute infectious mononucleosis. *N Engl J Med* 293:1159, 1975.
- Okano M, Purtilo DT: Interferons and Epstein-Barr virus infection. In Jaffe HS, Bucalo LR, Sherwin SA (eds): "Anti-Infective Applications of Interferon-Gamma." Basel, New York, and Hong Kong: Marcel Dekker, Inc., 1992, p 213.
- Okano M, Thiele GM, Kobayashi RH, Davis JR, Synovec MS, Grierson HL, Jaffe HS, Purtilo DT: Interferon-gamma in a family with X-linked lymphoproliferative syndrome with acute Epstein-Barr virus infection. *J Clin Immunol* 9:48, 1989.
- Imashuku S, Hibi S: Cytokines in haemophagocytic syndrome. *Br J Haematol* 77:438, 1991.
- Okano M, Thiele GM, Purtilo DT: Variable presence of circulating cytokines in patients with sporadic or X-linked lymphoproliferative disease with fatal infectious mononucleosis. *Pediatr Hematol Oncol* 10:97, 1993.
- Tomkinson BE, Wagner DK, Nelson DL, Sullivan JL: Activated lymphocytes during acute Epstein-Barr virus infection. *J Immunol* 139:3802, 1987.
- Akashi K, Hayashi S, Gondo H, Mizuno S, Harada M, Tamura K, Tamasaki K, Shibuya T, Uike N, Okamura T, Miyamoto T, Niho Y: Involvement of interferon-gamma and macrophage colony-stimulating factor in pathogenesis of haemophagocytic lymphohistiocytosis in adults. *Br J Haematol* 87:243, 1994.
- Kikuta H, Taguchi Y, Tomizawa K, Kojima K, Kawamura N, Ishizaka A, Sakiyama Y, Matsumoto S, Imai S, Kinoshita T, Koizumi S, Osato T, Kobayashi I, Hamada I, Hirai K: Epstein-Barr virus genome-positive T lymphocytes in a boy with chronic active EBV infection associated with Kawasaki-like disease. *Nature* 333:455, 1988.
- Mori M, Kurozumi H, Akagi K, Tanaka Y, Imai S, Osato T: Monoclonal proliferation of T cells containing Epstein-Barr virus in fatal infectious mononucleosis. *N Engl J Med* 327:58, 1992.
- Revoltella RP, Vigneti E, Fruscalzo A, Park M, Ragona G, Rocchi G, Calef E: Epstein-Barr viral DNA sequences in precursor monocyte-macrophage cell lines established from bone marrow of children with maturation defects of hematopoiesis. *J Gen Virol* 70:1203, 1989.
- Dolezal MV, Kamel OW, Rijn MVD, Cleary ML, Sibley RK, Warnke RA: Virus-associated hemophagocytic syndrome characterized by clonal Epstein-Barr virus genome. *Am J Clin Pathol* 103:189, 1995.
- Miyashita EM, Yang B, Lam KMC, Crawford DH, Thorley-Lauson DA: A novel form of Epstein-Barr virus latency in normal B cells in vivo. *Cell* 80:593, 1995.
- Okano M, Thiele GM, Gross TG, Davis JR, Purtilo DT: Differential cellular susceptibility to Epstein-Barr virus infection in a patient with X-linked lymphoproliferative disease. *J Med Virol* 32:47, 1990.
- Schuster V, Seidenspinner S, Kreth HW: Detection of a nuclear antigen 2 (EBNA2)-variant Epstein-Barr virus strain in two siblings with fatal lymphoproliferative disease. *J Med Virol* 48:114, 1996.
- Pracher E, Grumayer-Panzer ER, Zonbek A, Peters C, Gadner CP: Bone marrow transplantation during fulminant EBV-infection in Duncan's syndrome. *Lancet* 342:1362, 1993.
- Ambruso DR, Hays T, Zwartjes WJ, Tubergen DG, Favara BE: Success-

- ful treatment of lymphohistiocytic reticulosis with phagocytosis with epipodophyllotoxin VP16-213. *Cancer* 45:2516, 1980.
36. Chen R-L, Lin K-H, Lin D-T, Su I-J, Huang L-M, Lee P-I, Hseih K-H, Lin K-S, Lee C-Y: Immunomodulation treatment for childhood virus-associated hemophagocytic lymphohistiocytosis. *Br J Haematol* 89:282, 1995.
37. Oyama Y, Amano T, Hirakawa S, Hironaka K, Suzuki S, Ota Z: Hemophagocytic syndrome treated with cyclosporin A: A T cell disorder? *Br J Haematol* 73:276, 1989.
38. Fisher A, Blanche S, Bidois JL, Bordigoni P, Garnier JL, Niaudet P, Morinet F, Deist FL, Fisher AM, Griscelli C, Hirn M: Anti-B cell monoclonal antibodies in the treatment of severe B cell lymphoproliferative syndrome following the bone marrow and organ transplantation. *N Engl J Med* 324:1451, 1991.